

Cal/Ecotox

Exposure Factors for Northern Flicker (Colaptes auratus)*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Age at Sexual Maturity	review				NR	NR	CANADA; USA	a	1
Body Weight - Mean	140		131-144	g	NR	Adult	CO	b	2
Clutch or Litter Size	8			eggs/clutch	F	Adult	NC	c	3
Clutch or Litter Size	4.20			eggs/clutch	F	Adult	Lab	d	4
Clutch or Litter Size	6.5	1.4 SD	3-12	eggs/clutch	F	Adult	Lab	e	5
Clutch or Litter Size	6.18			eggs/clutch	F	Adult	OH	f	6
Clutch or Litter Size	4.71			eggs/clutch	F	Adult	OH	g	6
Clutch or Litter Size			6.97-7.75	eggs/clutch	F	Adult	NE; WY	h	7
Clutch or Litter Size			4-14	eggs/clutch	F	Adult	USA	i	8
Clutch or Litter Size	7.44	2.24 SD		eggs/clutch	F	Adult	OH	j	9
Dietary Composition	animal food (67.71%), vegetable food (32.26%)				NR	NR	CANADA; USA	k	10
Dietary Composition	predacious ground beetles (3.89%), other beetles (2.66%), ants (53.82%), Hemiptera (1.84%), caterpillars (2.12%), Orthoptera (1.45%), other insects and spiders (1.96%), grain (2.26%), seeds (19.59%), fruit (10.28%)				NR	NR	CANADA; USA	l	10
Dietary Composition	review				NR	NR	CANADA; USA	m	1
Duration of Incubation or Gestation	11			d	NR	Embryo	NC	n	3
Duration of Incubation or Gestation	review				NR	NR	CANADA; USA	o	1
Fledging or Weaning Rate	5.09			fledglings/c lutch	NR	Fledgling	OH	p	6
Fledging or Weaning Rate	2.75			fledglings/c lutch	NR	Fledgling	OH	q	6
Foraging Distance			0-1.4	km	NR	Adult	CO	r	2
Growth Rate	review				NR	NR	CANADA; USA	s	1
Hatching Success			0.826-0.903		B	Adult	NE; WY	t	7
Hatching Success	3.25			nestlings/clutch	NR	Nestling	OH	u	6
Hatching Success	5.27			nestlings/clutch	NR	Nestling	OH	v	6
Home Range			48-101	ha	NR	Adult	CO	w	2
Population Density	1.26 (20)			pairs/square mile	B	Adult	ND	x	11
Population Density	review				NR	NR	CANADA; USA	y	1
Time of Fledging or Metamorphosis	mid-June to late July				B	Adult	OH	z	9
Time of Fledging or Metamorphosis	review				NR	NR	CANADA; USA	aa	1
Time of Mating/ Laying	review				NR	NR	CANADA; USA	ab	1
Time of Mating/ Laying	review				NR	NR	CANADA; USA	ac	1
Time of Migration or Dispersal	review				NR	NR	CANADA; USA	ad	1
Time of Molt	June-July				NR	Both Adult and Juv.	Lab	ae	12
Time of Molt	review				NR	NR	CANADA; USA	af	1
Time of Nesting	early May to late June				B	Adult	OH	ag	9
Time of Nesting	mid-April to early May				B	Adult	OH	ah	9

Notes	
a	N=NR
b	average body weight at time of capture; N=4 birds; Feb-Apr; Lakewood, Jefferson County
c	first clutch size; N=1 nest; May; Asheville, Buncombe County
d	average size of clutches existing in North American oological collections; N=57 clutches; Clutch size for this subspecies is significantly smaller than for other subspecies.
e	average size of clutches existing in North American museum collections; N=411 clutches; Mean clutch size increases by ~1 egg per 10 degree increase in latitude.
f	average size of clutches laid early in the breeding season; N=17 clutches; May-June; New Concord, Muskingum County (40 deg, 00'N, 81 deg, 46'W)
g	average size of clutches laid late in the breeding season (1990-92,1994); N=7 clutches; May-June; New Concord, Muskingum County (40 deg, 00'N, 81 deg, 46'W); Late clutches were significantly smaller than early clutches.
h	range of mean clutch sizes measured among auratus ssp., cafer ssp., and auratus/cafer hybrid flickers; N=10-32 clutches per subspecies or hybrid group; Lincoln, Morrill, Scottsbluff Counties, NE; Platte County, WY; No statistical differences were found in clutch size among flicker subspecies or hybrids.
i	range of clutch sizes; N=191 clutches; The most frequently observed clutch sizes were 6, 7, and 8 eggs. Data are from eastern and southern USA.
j	N=9 pairs; spring; near New Concord; Clutch size was significantly smaller when European starlings were near nest.
k	percent composition of stomach contents; N=183 birds; all
l	percent composition of stomach contents; N=183 birds; all; Approximately 3/4 of sampled birds were collected in California. Species name given in report is <i>Colaptes cafer</i> . See paper for data on seasonal dietary changes and species lists of dietary items.
m	N=NR
n	incubation period; N=1 nest; May; Asheville, Buncombe County
o	N=NR
p	average number of fledglings from clutches laid early in the breeding season (1990-1992, 1994); N=11 clutches; May-June; New Concord, Muskingum County (40 deg, 00'N, 81 deg, 46'W)
q	average number of fledglings from clutches laid late in the breeding season (1990-1992, 1994); N=4 clutches; May-June; New Concord, Muskingum County (40 deg, 00'N, 81 deg, 46'W); Late clutches produced significantly fewer fledglings than early clutches.
r	range of distances between night roost location and the following day's locations; N=4 birds, 20-99 radio fixes per bird; Feb-Apr; Lakewood, Jefferson County
s	N=NR
t	range of mean brood to clutch size ratios measured among male and female auratus ssp., cafer ssp., and auratus/cafer hybrid flickers; N=4-42 clutches per subspecies or hybrid group; Lincoln, Morrill, Scottsbluff Counties, NE; Platte County, WY; No statistical differences were found in brood to clutch size ratio among flicker subspecies or hybrids.
u	average number of nestlings from clutches laid late in the breeding season (1990-1992, 1994); N=4 clutches; May-June; New Concord, Muskingum County (40 deg, 00'N, 81 deg, 46'W); Late clutches produced significantly fewer nestlings than early clutches.
v	average number of nestlings from clutches laid early in the breeding season (1990-1992, 1994); N=11 clutches; May-June; New Concord, Muskingum County (40 deg, 00'N, 81 deg, 46'W)
w	range of home range size based on telemetry location data; N=4 birds, 20-99 radio fixes per bird; Feb-Apr; Lakewood, Jefferson County
x	mean density of breeding birds (maximum density in parentheses); N=130 sample units (160 acres/unit); April-July
y	N=NR
z	N=40 pairs; spring; near New Concord; See paper for frequencies over time.
aa	N=NR
ab	N=NR
ac	N=NR
ad	N=NR
ae	date of onset of postjuvenal and adult molt; N=2500 birds; Data were based on examination of preserved specimens and live birds.
af	N=NR
ag	time of incubation; N=40 pairs; spring; near New Concord
ah	time of nest excavation starts; N=40 pairs; spring; near New Concord; See paper for frequencies over time.

References

1 Moore, William S. 1995. Northern Flicker (*Colaptes auratus*). In: Poole, A., P. Stettenheim and F. Gill, eds. The Birds of North America, No. 166. Philadelphia, PA: The Academy of Natural Sciences; Philadelphia, PA: The American Ornithologists' Union, Washington, D.C. p 1-28.

2 Royall, W.C., Jr., and O.E. Bray. 1980. A study of radio-equipped flickers. North Am. Bird Bander. 5(2):47-50.

3 Duyck, Bill E. and Douglas B. McNair. 1991. Notes on egg-laying, incubation, and nestling periods and of food brought to the nest by four species of cavity-nesting birds. Chat. 55(2):21-29.

4 Koenig, Walter D. 1984. Clutch size of the gilded flicker. Condor. 86:89-90.

5 Koenig, Walter D. 1984. Geographic variation in clutch size in the northern flicker (Colaptes auratus): support for Ashmole's hypothesis. Auk. 101:698-706.

6 Ingold, Danny J. 1996. Delayed nesting decreases reproductive success in northern flickers: implications for competition with European starlings. J. Field Ornithol. 67(2):321-326.

7 Moore, William S. and Walter D. Koenig. 1986. Comparative reproductive success of yellow-shafted, red-shafted and hybrid flickers across a hybrid zone. Auk. 103:42-51.

8 Burns, Frank L. 1900. A monograph of the flicker (Colaptes auratus). Wilson Bull. (31)1-82.

9 Ingold, Danny J. 1998. The influence of starlings on flicker reproduction when both naturally excavated cavities and artificial nest boxes are available. Wilson Bull. 110(2):218-225.

10 Beal, F.E.L. 1911. Food of the woodpeckers of the United States. Washington, D.C.: Government Printing Office. 7-64 p.

11 Stewart, Robert E., and Harold A. Kantrud. 1972. Population estimates of breeding birds in North Dakota. Auk. 89:766-788.

12 Test, Frederick H. 1945. Molt in flight feathers of flickers. Condor. 47:63-72.

*Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.